



King's Research Portal

DOI:

[10.1007/s00429-018-1676-3](https://doi.org/10.1007/s00429-018-1676-3)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Barttfeld, P., Abboud, S., Lagercrantz, H., Adén, U., Padilla, N., Edwards, A. D., Cohen, L., Sigman, M., Dehaene, S., & Dehaene-Lambertz, G. (2018). A lateral-to-mesial organization of human ventral visual cortex at birth. *Brain structure & function*. <https://doi.org/10.1007/s00429-018-1676-3>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

A lateral-to-mesial organization of human ventral visual cortex at birth.

Barttfeld P^{1,2*}, Abboud S^{4,5*}, Lagercrantz H⁶, Adén U⁶, Padilla N⁶., Edwards AD⁷, Cohen L^{4,5}, Sigman M³, Dehaene S^{1,8} & Dehaene-Lambertz G¹

- 1- Cognitive Neuroimaging Unit, CEA DSV/I2BM, INSERM, Université Paris-Sud, Université Paris-Saclay, NeuroSpin Center, Gif-sur-Yvette, France
- 2- Laboratory of Experimental Psychology and Neuroscience (LPEN), INECO (Institute of Cognitive Neurology)
- 3- Universidad Torcuato Di Tella, Almirante Juan Saenz Valiente 1010, C1428BIJ Buenos Aires, Argentina.
- 4- INSERM, U 1127, Paris, F-75013, France.
- 5- Institut Du Cerveau Et De La Moelle Epinière, ICM, PICNIC Lab, Paris F-75013, France.
- 6- Department of Women's and Children's Health, Karolinska Institutet, 171 76 Stockholm , Sweden.
- 7- Centre for the Developing Brain, Division of Imaging Sciences & Biomedical Engineering, King's College London, London, UK.
- 8- Collège de France, 75005 Paris, France.

*These authors contributed equally to the work.

Abstract

In human adults, ventral extra-striate visual cortex contains a mosaic of functionally specialized areas, some responding preferentially to natural visual categories such as faces (fusiform face area) or places (parahippocampal place area) and others to cultural inventions such as written words and numbers (visual word form and number form areas). It has been hypothesized that this mosaic arises from innate biases in cortico-cortical connectivity. We tested this hypothesis by examining functional resting-state correlation at birth using fMRI data from full-term human newborns. The results revealed that ventral visual regions are functionally connected with their contra-lateral homologous regions and also exhibit distinct patterns of long-distance functional correlation with anterior associative regions. A mesial-to-lateral organization was observed, with the signal of the more lateral regions, including the sites of visual word and number form areas, exhibiting higher correlations with voxels of the prefrontal, inferior parietal and temporal cortices, including language areas. Finally, we observed hemispheric asymmetries in the functional correlation of key areas of the language network that may influence later adult hemispheric lateralization. We suggest that long-distance circuits present at birth constrain the subsequent functional differentiation of the ventral visual cortex.

Keywords

Neonates – functional connectivity - language

Introduction

The human brain displays a remarkable consistency and reproducibility in its functional organization across individuals. This was first demonstrated in neuropsychological studies that mapped specific cognitive deficits to reproducible cortical lesion sites and more recently by brain-imaging research, which has revealed surprisingly similar functional subdivisions of the cortical surface across numerous individuals (Glasser et al., 2016; Huth, de Heer, Griffiths, Theunissen, & Gallant, 2016). Not only is this reproducible topological specialization evident in domains of evolutionary relevance, such as for the fusiform face area (FFA) and the parahippocampal place area (PPA) (Downing et al., 2006) but it also extends to cultural acquisitions such as reading. For example, the same region of the left fusiform gyrus, the visual form area (vWFA), is consistently activated whenever literate participants recognize written words, independently of case (Dehaene & Cohen, 2011), modality (Striem-Amit, Cohen, Dehaene, & Amedi, 2012) or the age of reading acquisition (Dehaene et al., 2010).

Such functional reproducibility suggests that cognitive processes are dependent on specific neural architectures (Hannagan, Amedi, Cohen, Dehaene-Lambertz, & Dehaene, 2015). Indeed, functional responses have been consistently related to specific micro-architectures (e.g. word and face-specific activations are observed in the regions FG2 and FG4, whereas places are activating FG1 and FG3, Caspers et al., 2015; Weiner et al., 2016), although the exact relation between a given cortical layer organization and a computational process are still poorly specified. Anatomical connectivity, by orchestrating local and distant synchrony, might also drive the functional role of an area. This “biased connectivity” hypothesis was recently supported by two studies showing that the precise locations of the FFA (Saygin et al., 2012) and VWFA (Saygin et al., 2016) can be predicted from their connections. The functional localization of the VWFA was predicted in 8-year-old readers by the anatomical fiber tracking performed at the age of 5 before they learned to read (Saygin et al., 2016).

Because all main fiber tracts are already present in full-term human neonates (Dubois et al., 2014; Dubois et al., 2016), in the present paper, we asked whether the initial connectome, present at birth, may explain at least some aspects of the subsequent functional specialization. We focused here on the mesial-to-lateral organization of ventral visual cortex. The ventral visual pathway comprises a succession of functional areas extending from mesial to lateral regions including the mesial place area (PPA), the face fusiform area (FFA), the visual word form area (VWFA), the number form area (NFA) and a more lateral area responding to tools and objects. Furthermore, distinct face and place responses are already present in 6-month-old infants (Deen et al, 2017). Here, we examined whether a particular connectivity pattern of these regions is already in place at birth.

To this aim, we relied on resting-state functional correlation, i.e. the synchronization of the BOLD responses across distant regions at rest due to the spontaneous co-activity of different nodes of the same network (Glasser et al., 2016; Raichle, 2015; Smith et al., 2013). Using this approach in full-term and even pre-term neonates, several studies have reported the presence of the main resting-state networks described in adults (Doria et al., 2010; Fransson, Aden, Blennow, & Lagercrantz, 2011; Fransson et al., 2009; Fransson et al., 2007; Smyser et al., 2010; Smyser, Snyder, & Neil, 2011). Our goal was to go beyond the whole-brain description and determine whether long-distance connections specifically arising from ventral visual areas already target distinct regions. We aimed to evaluate two contrasting theories for the origins of adult functional specialization. The first of which, the “biased connectivity” hypothesis (Hannagan et al., 2015; Saygin et al., 2012) as explained above, predicts that mesial-to-lateral differences in functional correlation are already present at birth, in approximate correspondence to the functional mosaic later found in older children and adults. We then contrasted it to the latter hypothesis according to which brain connectivity is initially diffuse, over-abundant and largely isotropic and only becomes refined at a later stage during infancy and childhood through exposure to a given environment (Elman et al., 1996; Johnson, 2011).

The “biased connectivity” hypothesis makes specific predictions for which we searched in our data. Given the reproducibility of the location of the visual word and number form areas, we wondered whether these regions were already more connected with the temporo-frontal language network and the inferior parietal “number sense” regions than other mesial regions, which would explain their propensity to acquire the role of processing arbitrary shapes to represent phonemes or numbers. Similarly, we wondered whether hemispheric biases in connectivity at birth already predated the later lateralized responses to faces and words in the right and left fusiform gyrus (Weiner et al, 2016). We thus analyzed whether key regions of the linguistic network (i.e. pars triangularis of the inferior frontal gyrus, superior and middle temporal gyrus) are differently related to the left and right ventral temporal regions.

Methods

1. Subjects

Twenty-four healthy full-term neonates scanned for research purposes were included in this study. The fMRI data were provided by the Karolinska University Hospital, Stockholm (14 neonates) and by the Centre for the Developing Brain, King’s College London (10 neonates) and have already been published (Fransson et al, 2009; Doria et al, 2010). All MR examinations were carried out according to the ethical guidelines and declarations of the Declaration of Helsinki (1975). Details can be found in the seminal papers. Six subjects were subsequently rejected due to excessive movement artefacts (see preprocessing section). We thus analyzed XX girls and XX boys aged XX days in average at scan.

2. MR Image Acquisition

For the Swedish cohort, infants were scanned during natural sleep. Functional MRI scans were acquired on a Philips Intera 1.5 Tesla scanner equipped with a 6-channel receive-only head coil. Echo planar imaging (time repetition [TR]/time echo [TE]/flip angle = 2000 ms/50ms/80°,

voxel size=2.8 x 2.8 x 4.5 mm, 20 slices) of the infant brain was performed for 10 minutes (300 EPI volumes). For the London cohort, infants were sedated with chloral hydrate (25–50 mg/kg) and images were acquired at 3 Tesla (Achieva, Philips, Best, Netherlands) with an 8 channel phased array head coil. Echo planar imaging (TR/TE/flip angle = 1500ms/45ms/90°, 2.5 x 2.5 x 3.25mm, 22 slices) was performed for 6.4 minutes. In both groups, T1-weighted images were acquired in each neonate. Although it is unconventional to merge data obtained from two different MRI scanners for combined group-level inference, we were confronted to the scarcity of the publically available data at this age and the small size of the groups. Note that our first-level comparisons were computed within-subject searching for differences in functional connectivity between voxels. By merging the two groups, we probably increased the variance between subjects but we benefited from more degrees of freedom. To capture the group differences, a group-factor was declared in our statistical analyses.

3. Preprocessing.

FMRI images were all pre-processed using SPM8 (Wellcome Institute of Imaging Neuroscience, London, UK) implemented in Matlab (Mathworks, Natick, MA). Pre-processing involved within-subject realignment to account for head motion, spatial normalization to a template and smoothing: all images were realigned to the mean image of the resting state session using linear transformations. The mean EPI image was spatially normalized to a 7 week-old infant template image for whom a sulci-based parcellation is available (Kabdebon et al., 2014), thus providing anatomically defined region of interests to study functional connectivity. Brain growth and sulcation massively develop during the last trimester of gestation but decelerate after term-age. Although tertiary sulcation begins around full-term birth and continues to develop until adulthood, all main structures important for normalization are visible (see Kabdebon et al, 2014, for quantitative measures of inter-age variability of the location of the main sulci during the first semester of life). The transformation parameters obtained from this spatial normalization were subsequently applied to every functional image acquired during the resting state scanning session

with a target voxel size of $2 \times 2 \times 2 \text{ mm}^3$. Then, spatially normalized images were smoothed with an isotropic 5-mm^3 full-width half-maximum Gaussian kernel. Next, we regressed out the effect of the following nuisance variables from the voxel time series using the REST toolbox for Matlab (<http://restfmri.net>): the six movement parameters from the head motion correction stage along with the global signal of the entire brain and four nuisance signals obtained from a 3-mm sphere located in the ventricles and white matter. Finally, all functional MRI time series were band-pass filtered (0.01 - 0.08 Hz).

Spontaneous movement during sleep in infants often consists of sudden hyperextensions along the body axis which produced spiking artefacts in the time series and might have therefore disrupted anterior-posterior correlations as a result of these pitch movements. Because subject motion is particularly deleterious on long-distance correlations (Power, Schlaggar, & Petersen, 2015), we removed the artifacted time points through a semi-automatic method. First, we deleted brain volumes detected by the ART toolbox for SPM8 as contaminated by high levels of movement. Second, we visually inspected whether all traces of movement were removed from the voxel time series. When needed, more brain volumes were discarded until no trace of movement was visually detected in the time series. Concerns about such scrubbing have been raised in the literature as it decreases the degrees of freedom and destroys the temporal time-series when too many successive brain volumes are removed (for a discussion see Power et al., 2015). Furthermore, given the non-stationarity of the functional correlations, some stages during which subjects might move more may be under-represented. These concerns are particularly critical when different populations with different levels of agitation are compared, such as infants with adults. However, this concern does not apply to our study, since we are comparing the connectivity between different voxels that have been similarly processed within subjects. In the end, six subjects whose remaining time-series were shorter than 130 scans were discarded to avoid noisy correlation estimations. All included subjects had at least 138 brain volumes (mean number of volumes: 218, SD: 47, min: 132, max: 300).

4. Clustering analysis

To parcel the ventral areas in clusters of voxels with similar patterns of functional correlation, we applied the following data-driven clustering method. First, we performed a functional correlation analysis (Brain Voyager, Neuroelf v1.0, <http://neuroelf.net/>) seeded from each voxel belonging to a mask of the ventral stream of the visual cortex, i.e., ventrally to the Calcarine sulcus expanding towards the anterior parahippocampal gyrus and limited to $y = -14$ in the template (1484 voxels, corresponding to the union of the clusters presented in figure 1). In other words, in each individual neonate, time-series were extracted from each functional voxel within the mask z-normalized and used as general linear model (GLM) predictors. For each voxel in the mask, random-effects inference was then performed on the group level. Second, each resulting group-level map was masked by a gray-matter mask, thresholded (voxelwise $p < 0.01$ uncorrected), binarized and reshaped into a numerical vector. We then represented the functional correlation from all seeds as a matrix of 1484 columns (the number of seeds) and 14808 rows (the number of gray-matter brain voxels). Third, we used an agglomerative hierarchical clustering algorithm to cluster the seed voxels with a similar spatial pattern of functional correlation (i.e. a similar matrix columns). A hierarchical cluster tree was constructed using the Ward linkage measure on Euclidean distances (minimum variance algorithm merging at each step the pair of clusters that leads to the minimal increase in total within-cluster variance). The final number of clusters was optimized using the silhouette technique which quantifies the ratio between the largest inter-cluster distance over the smallest distance to a point outside the cluster. An assessment of all solutions in the range of 2 to 15 clusters showed a local maximum for 4 clusters. Fourth, in order to study the brain networks correlated with each of these 4 clusters, we performed a group-level analysis similar to the one described above, but using the average time-series from all the voxels of each cluster computed in each individual neonate. We also computed partial correlation coefficients for each cluster in order to regress out the time-series of all other clusters and thus obtain its unique correlation pattern (figure 1). Significance thresholds were set to $q = 0.01$ (FDR-corrected for multiple comparisons)

when using partial correlation and $q=0.001$ (FDR-corrected for multiple comparisons) when using Pearson's correlation.

5. Seed analysis

Using the anatomical parcellation proposed by Kabdebon et al (2014) on a two-month-old infant template, we defined four ROIs based on the gray matter of 4 sulci bordering the lingual, fusiform (internal part), fusiform (external part), and the inferior temporal gyrus. Those ROIs were used as seeds for the correlation analysis (figure 2a). For each subject and each ROI, we defined the time-series T as the average time-series across all the voxels within the ROI. Then, we calculated the Pearson correlation between T and the time-series of all the voxels in the brain, V . Through this procedure, we obtained 4 correlation brain volumes or maps per subject s and ROI r ($C_{s,r}$), which were Fisher-transformed (Matlab function `atanh`) to produce $Z_{s,r}$. To test for significant correlations across subjects between T and all voxels in the brain, we performed a one-way t-test on each ROI r , setting the significance value at $p < 0.01$, FDR-corrected voxelwise. We also performed a within-subjects ANOVA between maps so as to compare pairs of $Z_{s,r}$ for each individual subject, setting the significance value at $p < 0.01$ for visualization purposes (a more strict threshold of $p < 0.001$ did not qualitatively change the results).

6. Regression analysis

Given that the boundaries between ROIs in the previous analysis were based on sulci that might not follow cytoarchitectonic boundaries (Fischl et al., 2008), we performed an analysis free of this constraint through voxel-wise regression to explore the mesial-to-lateral gradient in a ROI-independent manner. We constructed a mask comprising both left and right ventral regions (Figure 3a). We reduced this mask to a x,y grid along the left-right and antero-posterior axes, merging together all voxels along the z axis (dorsal-ventral axis), as the mosaic of activations to visual categories that we wanted to explore is mainly evident along these two axes (Weiner et al, 2016). We computed the correlation map for each x,y pair and the time series of all voxels in the brain in

each subject and obtained a total of 276 (for left hemisphere) and 258 (for right hemisphere) correlation maps per subject. To identify regions whose correlation with voxels in the ROI would depend on their x and y coordinates, we performed a first-level regression analysis (using SPM) for each subject. We then submitted the x-related contrast to a second-level analysis, in order to quantify the influence of the x coordinate on functional correlation. We modeled the cohort origin (i.e. England or Sweden) explicitly as blocks in the analysis. The resulting second-level beta map and its associated t-map revealed the relation between the x-axis and functional correlation: the higher the beta and the t value, the more connected each voxel was with increasing values of x (i.e. from medial to lateral localization). Conversely, the smaller the effect the more connected each voxel was with decreasing values of x. We performed this procedure for the left and the right hemispheres, inverting the x values when necessary to keep the convention that increasing x values correspond with lateral locations.

We wondered whether the differences we observed between mesial and lateral regions were related to their level of maturation which might have created different delays in neurovascular coupling. As a control procedure aimed at accounting for this possibility, we repeated the analysis including the average T2 value of the template brain at the same locations as a regressor and thus a proxy for maturation index (Leroy et al., 2011).

7. Analysis of hemispheric differences

Because the VWFA and the FFA are respectively left and right lateralized in adults, we studied whether differences in functional correlation were observed early in development, between the ventral areas and key regions of the linguistic network. Therefore, we recovered the following anatomically pre-defined ROIs from the infant atlas provided by Kabdebon et al. (2014): the pars triangularis of the inferior frontal gyrus (region number 4 in the atlas), the middle temporal gyrus (region 27) and the superior temporal gyrus (region 28). We truncated regions 27 and 28 along the anterior-posterior axis at the level of the deepest angle of the *planum temporale* so as to keep the

ROIs most anterior parts. This was done to avoid any overlap or vicinity of the seeds with the visual stream that could have biased the analysis with spurious local correlations.

For every subject and every pair of left-right regions, (e.g. left and right pars triangularis), we performed two seed analyses, the first between the left ROI time-series and the rest of the brain and the second using the right ROI time-series. We thus obtained two functional correlation maps per subject and ROI pair. To test for hemispheric laterality biases, we performed a series of t-test (for each value of x coordinate) on the left-right difference collapsing all y-coordinate values between values of y coordinate of 9 and 13. The significance threshold was set at $p < 0.05$, Bonferroni corrected for multiple comparisons (42 comparisons, corresponding to the 42 voxels along the x coordinate).

Results

1. Parcellation of visual ventral regions

We first performed a data-driven clustering of the voxels in the bilateral ventral regions according to their functional correlation pattern. Voxels were clustered together or separately as a function of the similarity of the pattern of correlation with the entire brain. This clustering procedure was combined with a partial correlation analysis in order to display the unique functional fingerprint of each cluster (i.e. not shared with other clusters). We identified four main clusters: C1, or Medial Occipital, posterior, an exclusively posterior occipital group connected with the frontal pole, the right inferior temporal gyrus and paracentral regions consisting of the right post-central gyrus, paracentral lobule and supplementary motor area; C2, or Medial Occipital, anterior, connected with neighboring mesial areas (lingual, fusiform, parahippocampal gyri, and the cuneus) and more distant primary regions, such as the central sulcus, Heschl's gyrus, *planum temporale*, bilateral posterior insula and the center of the right STAP region (the STAP is a region under Heschl's gyrus in which the right superior temporal sulcus is deeper than the left in almost all humans, Leroy et al., 2015); C3, or Anterior Lateral, a cluster which had the most extended unique

functional correlation, connected to the inferior and middle temporal lobes, the inferior parietal regions comprising the inter-parietal sulcus, and the precentral gyri; C4, or Posterior Lateral, a cluster connected to the dorsal visual stream, cuneus and left dorso-lateral prefrontal region (Figure 1 and S1 for functional correlation using Pearson's correlation).

2. Functional correlation based on anatomical ROIs

The parcels defined above correspond to an optimal statistical solution given our data but may differ from the anatomical reality. We thus sought to confirm the above conclusions using four anatomical ROIs progressively aligned along the medial-external axis of the left ventral temporal cortex in order to determine how they connected to the rest of the brain. As expected from the previous analysis, the observed functional correlation was not limited to neighboring ipsilateral areas or to homologous regions in the right hemisphere (figure 2). Rather, the connections crossed hemispheres to join distant associative areas in the right hemisphere. We observed a different correlation pattern depending on ROI position (Figure 2b): the most mesial ROI (ROI1, or Lingual gyrus) functionally related to the superior temporal region as well as central motor and sensory regions whereas the most lateral ROI (ROI4) was principally correlated with the whole temporal lobe, the inferior parietal region and the prefrontal region. ROIs 1 and 4 thus shared similarities with the clusters C2 and C3 of the previous analysis.

Pairwise comparisons between the patterns of functional correlation of the different ROIs (figure 2c) revealed that ROI1 was most strongly correlated with the lower part of the central sulcus and Heschl's gyrus bilaterally. ROI2, the medial part of the Fusiform gyrus presented only local correlations which extended more anteriorly in the left fusiform and inferior temporal gyri relative to ROI1. A sharp change occurred with ROI3, the more lateral part of the Fusiform gyrus, which displayed correlations with the ventral regions in both hemispheres and moreover with the supra-marginal and angular gyri, also in both hemispheres. This change corresponds to the internal and external part of the fusiform gyrus, marked by the mid-fusiform sulcus (Weiner et al., 2014).

Finally, ROI4 displayed the most extended correlation pattern, exhibiting stronger correlations with the whole temporal lobe and prefrontal regions relatively to the other ROIs and notably to ROI3. It is noticeable that the significant changes of functional correlation patterns between ROI3 and ROI4 (The posterior part of the Temporal gyrus) spared the parietal and frontal regions.

A possible confound is a similar neurovascular coupling in regions at the same maturational stage. While there is no indication that equally maturing regions will show a correlated functional activity in time, to account for this possibility we repeated the analysis including the average T2 value of the template brain at the same location as a regressor and thus as a proxy for maturation index (Leroy et al., 2011). Because dendritic development impairs water motion in the gray matter and myelination in the white matter, T2 signal becomes progressively darker during the first year of life and can hence be used as a proxy for neural maturation (Leroy et al., 2011). We observed virtually the same functional correlation pattern, suggesting that maps presented in figure 2b, c are not directly related to a maturational neuro-vascular gradient.

A third, voxel-wise, analysis aimed to verify that the mesial-to-lateral organization of the fusiform gyrus was not biased by our arbitrary ROI definition. To this aim, we examined the functional correlation between each x,y location in a ventral region encompassing the calcarine sulci, lingual gyrus, fusiform gyrus and inferior temporal gyrus and examined the dependency of this functional correlation pattern on the x-coordinate of the voxel of origin (see Methods). In agreement with our previous ROI analysis, functional correlation was sensitive to the x-coordinates (figure 3): Heschl's gyrus and posterior insula signal showed a greater functional correlation with mesial voxels in the ventral visual cortex (thus appearing as a negative correlation in figure 3b), whereas associative areas of the temporal lobe (sparing the superior temporal region) in both hemispheres, the left angular and supramarginal gyri and both prefrontal regions showed a greater functional correlation with lateral voxels (thus appearing as a positive correlation in figure 3b; see Table S2).

3. Hemispheric asymmetries

Finally, to study putative asymmetries in the manner in which right and left ventral streams relate to the language network, we performed a within-subject comparison of the functional correlation maps arising from three pairs of symmetrical anatomical ROIs located at key language-related areas: pars triangularis in the inferior frontal gyrus (IFG), superior and middle temporal gyri (STG and MTG). Figures 3c-3k thus show to what extent each site in the left and right ventral visual cortex was more strongly correlated to the left or right IFG, STG and MTG. Unsurprisingly, STG and MTG exhibited a stronger correlation with the ipsilateral than with the contralateral hemisphere, but this relation was stronger in the left hemisphere at a peak close to the location of the VWFA in adults ($x=-36$ and $y=-30$, MNI coordinates). After correction for multiple comparisons, this peak remained significant only for the MTG (Figure 2 g, j). More surprisingly, the left IFG signal was more correlated than the right IFG with a right lateral ventral spot at $x=24$ and $y=-30$ (MNI coordinates).

Discussion

We explored the pattern of functional correlations of the ventral visual cortex with the rest of the brain in sleeping full-term neonates. Both of our analyses, whether data-driven (figure 1) or based on anatomically defined regions (figure 2), revealed a clear partition of the ventral regions supporting the hypothesis of an early functional organization. Mesial ventral areas were notably correlated with other primary regions whereas the most lateral regions were more correlated with the whole temporal lobe, the inferior parietal region and the prefrontal region with a transition occurring between ROI 2 and 3 (figure 2). This boundary roughly corresponds to the mid-fusiform sulcus which splits the fusiform gyrus (Weiner et al, 2014). This sulcus has been proposed as a robust cyto-architectonic landmark between FG1 and FG3 on one side and FG2 and FG4 on the other side, and functional landmark between the place area and the FFA/VWFA in adults (Weiner et al., 2016; Weiner et al., 2014). Finally, studying hemispheric differences in functional correlations,

two results emerged: (1) when comparing intra-hemispheric correlations, we observed a stronger correlation between the middle temporal gyrus and the fusiform region in the left than in the right hemisphere, and (2) for interhemispheric correlations, we observed a stronger correlation between a right lateral ventral area and the contralateral left IFG than with the ipsilateral right IFG.

This organization of the associative visual regions is all the more remarkable given that vision is very limited *in utero*, contrary to the other senses which are regularly stimulated through the fetus' spontaneous movements, internal and external sounds crossing the maternal abdomen, and a tasty and odorant amniotic liquid reflecting the mother's meals. This early organization of the visual areas at birth, beyond primary areas, dismisses the old but persisting idea of a blank slate newborn (Elman et al., 1996; Johnson, 2011). Although the human infant appears immature at birth due to his poor motor repertoire, early learning is thus supported by a pre-existing cortical architecture now extensively documented by a decade of functional brain imaging studies in infants (Arichi et al., 2010; Bartocci, Bergqvist, Lagercrantz, & Anand, 2006; Mahmoudzadeh et al., 2013; Mahmoudzadeh, Wallois, Kongolo, Goudjil, & Dehaene-Lambertz, 2016).

Our working hypothesis went beyond a mere organization at birth, and we studied whether the functional specificity described at later ages in ventral temporal areas for visual categories such as words, faces, tools and places might partly be based on a pre-existing pattern of functional correlation of these regions with the rest of the brain ("biased connectivity" hypothesis). Anatomically, long-range pathways, in particular those connecting frontal regions with temporal and occipital regions can be observed in post-mortem brains from the second trimester of gestation on (Takahashi, Folkerth, Galaburda, & Grant, 2011). Even if the tracts are initially not well myelinated, the structural architecture of the human brain set up during the last month of gestation offers the possibility of long-distance synchrony. Previous studies of functional correlations in full-term and preterm neonates have indeed reported a set of maps of functionally synchronized regions similar to those observed in adults (Fransson et al., 2007, 2009, 2011; Doria et al., 2010; Smyser et al., 2010, 2011). Here, we explored whether a finer-grained pattern of functional correlation was

present within ventral visual cortex, correlating with the functional specificity for visual categories observed in adults.

Our results indicate that the most lateral regions are more connected to distal associative regions, notably in the frontal lobe, than mesial regions are (compare cluster 3 and ROI 4 to cluster 2 and ROI 1 in figures 1 and 2, respectively). The privileged connectivity of such lateral regions residing at the top of the hierarchy of early visual regions may thus explain why the cultural symbols of letters and digits later develop at this location. This aspect of our results corroborate prior findings in older children, which have indicated that anatomical connectivity prior to reading acquisition (at 5 years of age) of ventral visual voxels to temporal and frontal regions was predictive of those voxels which would later exhibit a preference for written words at 8 years of age, once children had learned to read (Saygin et al., 2016). In both cases, connectivity (anatomical or functional) precedes functionality, thus supporting the “biased connectivity” hypothesis for the origins of ventral visual specialization.

By contrast, we found that mesial areas are functionally more connected to sensorimotor areas, Heschl’s gyrus and the posterior insula. This observation could be mediated by direct connections between visual primary areas and auditory and parietal cortices which have been reported in newborn kittens (Innocenti, Berbel, & Clarke, 1988) and adult macaques (Rockland & Ojima, 2003). Alternatively, the observed functional correlation between mesial visual regions and other primary cortices (auditory areas and motor sensory areas at the level of the mouth and tongue) could also be driven by thalamic synchrony. Alcauter et al (2014), studying thalamo-cortical connectivity during infancy, reported an overlap of the connectivity between primary visual, auditory and sensory-cortices within the same thalamus cluster in neonates, whereas distinct clusters were observed at one year of age. Contrarily, Toulmin et al (2015) described a more segregated thalamo-cortical connectivity from birth on, especially concerning the primary cortices. We may speculate that an initial stage of multisensory integration might facilitate learning by enhancing responses to co-occurring cues that need to be integrated into higher-level

representations. For example, it might explain the observations that neonates visually recognize their mother only if they have been simultaneously exposed to her face and voice (Sai, 2005) and that 3-month-olds more robustly identify their mother vs. a stranger's face if her voice is simultaneously presented (Burnham, 1993).

This mesial-lateral division of functional correlation was similarly observed in the left and right hemisphere (see figure 3b) and may correspond to the cyto-architectonic division highlighted in adults by Weiner et al. (2014, 2016), and to the functional parcellation of activations revealed in older infants looking to faces/objects vs. places images (Deen et al., 2017). In both hemispheres, long-distance functional correlation was not limited to within-hemispheric areas, but also recruited contra-lateral heterotopic areas encompassing the inferior parietal and dorsal frontal regions. Although roughly similar, interesting differences were observed between the two hemispheres. Because the VWFA is systematically lateralized to the hemisphere involved in spoken language processing (Cai, Paulignan, Brysbaert, Ibarrola, & Nazir, 2010; Pinel & Dehaene, 2009; Pinel et al., 2014), which is usually in the left hemisphere, we examined the functional correlation at birth between three key regions of the spoken language network and ventral visual areas (figure 3c-i). The STG and MTG were strongly connected with both the ipsi- and contra-lateral visual regions, but the left MTG was significantly more related with the left lateral extra-striate visual region than its right counterpart. Thus, a stronger path already exists at birth between the spoken language network and the future location of the VWFA. It may explain why reading acquisition always recruits this particular location within the left ventral visual cortex (Hannagan, Amedi, Cohen, Dehaene-Lambertz, & Dehaene, 2015).

The significant trans-hemispheric correlation between the left inferior frontal region and the right lateral ventral visual cortex was more surprising. This cluster is more mesial than the cluster discussed above (~24 vs -38 mm in our mask x-coordinates). Contrary to temporal regions, left and right inferior frontal regions were weakly correlated with ventral voxels in our mask except for this particular relation between this right cluster and the left frontal region. Note that in figure 3b, in

which we studied the functional correlation of the left and right ventral temporal regions, we also observed a crossed relation yet in the reverse direction between the left temporal region and the right inferior frontal region. Does this functional correlation pattern correspond to a genuine, anatomically validated pathway? In rare patients with a developmental agenesis of the corpus callosum, sigmoid bundles connecting the right frontal region with the left contra-lateral occipital region (and, even more rarely, the converse pattern) have been detected by diffusion-tensor imaging (Benezit et al., 2015; Tovar-Moll et al., 2007). On one hand, these cases may correspond to aberrant bundles. On the other hand, the absence of a corpus callosum may reveal a smaller tract, otherwise masked by the large corpus callosum. Indeed, in a post-mortem tracing study in a single adult human brain, Virgilio and Clarke (1997) were able to identify the degenerating axons in a patient who died a few weeks after a stroke in the right inferior temporal region and reported inter-hemispheric mono-synaptic connections between the right inferior temporal region and the left-hemispheric Wernicke's and Broca's areas. Thus, interhemispheric connections can be heterotopic and relate widely different regions, in agreement with the present results. Our functional correlation pattern may also be related to a rare study of face processing in infancy by Tzourio-Mazoyer and collaborators (2002). Using positron emission tomography, these authors reported activations in left inferior frontal and left superior temporal cortex, together with right fusiform cortex, when two-month olds were looking at faces. Efficient connections between right-hemispheric fusiform face representations and left-hemispheric language areas might explain why vocal imitations can be so easily elicited when infants view articulating faces (Bristow et al., 2009; Chen, Striano, & Rakoczy, 2004; Meltzoff & Moore, 1977).

Finally, the present results speak to major theories on the origins of functional specialization in the adult human brain. They show that the neonate connectivity landscape is very far from a "blooming buzzing confusion" (James, 1890). As such, our results refute theories of an extreme empiricist nature, which assumes that the infant brain is largely devoid of initial architecture and that connectivity is initially diffuse, over-abundant and largely isotropic, and only becomes refined

at a later stage during infancy and childhood through exposure to a given environment (Elman et al., 1996; Johnson, 2011; Quartz & Sejnowski, 1997). Ventral visual cortex is not just “trained” by exposure to objects, faces, houses, letters or digits; on the contrary, it contains functionally specialized areas already present at the dawn of infancy (Biagi, Crespi, Tosetti, & Morrone, 2015; Deen et al., 2017), supported by well-defined fiber tracts. Several previous studies have already shown how at a global scale, an organized functional and anatomical connectivity is present in the infant brain (Fransson et al., 2007, 2009, 2011; Doria et al., 2010; Smyser et al., 2010, 2011; van den Heuvel et al., 2015). Here we show a refined organization within a single region with a gradient of preferred relations with local and distant regions, in accordance with the hypothesis of an early organization (“protomaps”) determined during gestation (Mahmoudzadeh et al., 2016; Rakic, 1988; Toulmin et al., 2015). These maps provide neonates with a blueprint on which learning capitalizes.

While indisputable differences in functional correlation were found between mesial and lateral regions, we did not recover a full adult-like organization. For instance, given the systematic location of word- and number-form areas in adults (VWFA and NFA), with the NFA systematically occupying a location lateral to the VWFA (Amalric & Dehaene, 2016; Hannagan et al., 2015; Shum et al., 2013), we could have expected a differential pattern of connectivity, with the most lateral ventral visual voxels projecting preferentially to the inferior parietal lobe which has been implicated in number sense; however, such was not the case. That being said, our method was crude in the sense that we did not know precisely which cortical sectors, in our infants, would later become specialized for letters versus digits, unlike what had been done in older children (Saygin et al., 2016). Future studies with an improved methodology might reveal that the infant brain contains even finer-grained biases that predate the fine differences between VWFA and NFA seen in adults. Alternatively, it is possible that such patterns of functional specialization emerge through exposure and learning and that initially only a rough division between mesial and lateral regions is visible. In line with this idea, Deen et al (2017) reported clear differences between fMRI activations for faces vs. places but not for objects vs. faces in 4-6 month-old infants. Similarly, Gomez et al. (2017)

underlined the protracted development of the fusiform region sensitive to face relative to more mesial areas. The acquisition of longitudinal data in the same children from infancy to childhood, although extremely difficult, may be necessary to answer this question.

Acknowledgements

Authors would like to thank Elvis Dohmatob, Isabelle Denghien, François Leroy, Jessica Dubois for their constructive remarks during this work.

Figure Legends

Figure 1. **Clustering of ventral visual cortex according to long-distance whole-brain functional correlations in newborns.** a) Four clusters were identified and are presented by different colors on axial slices of the infant template. b) Partial correlation maps showing the distinct patterns of whole-brain correlations associated with each of the four clusters. All clusters exhibit long-distance functional correlations, but the most lateral and anterior cluster (C3) appears to be specifically correlated to inferior parietal and frontal associative areas.

Figure 2. **A mesial-lateral gradient of functional correlation in ventral visual cortex.** a) The four ROIs used in this analysis as seeds, ordered from mesial to lateral. b) Functional correlations of the 4 ROIs, rendered on a 3D infant inflated brain. $p = 0.01$, FDR corrected. c) The connectivity patterns of each pair of ROIs are compared two by two. Each column presents the t-contrast row ROI > column ROI. (For illustration purposes, p-value was set at $p < 0.01$, uncorrected.) A mesial-to-lateral gradient is clearly visible: mesial seeds preferably correlate with primary regions, while lateral ROIs are more strongly related to associative regions in the temporal, parietal and frontal lobes.

Figure 3. **Hemispheric differences in long-distance functional correlations.** a) The mask containing all voxels used as seeds in the regression analysis is presented on a horizontal slice. To quantify to what extent functional correlations vary with the x-coordinate, a connectivity map was computed for each voxel in this mask and was subsequently entered into a regression analysis with the x coordinate as a regressor. b) Functional correlation pattern as a function of the x-axis location of the seed in the left and right hemispheres. The more positive the value, the stronger the correlation with lateral voxels (e.g. the right inferior frontal region with a lateral left seed) whereas the more negative the value the stronger the correlation with mesial voxels (e.g. Heschl's gyrus on both sides). c-k) Analyses of hemispheric biases. c, f, i display the location of the three bilateral ROIs used as seeds: inferior frontal gyrus (c), medial temporal gyrus (f) and superior temporal

gyrus (i). d, g, j display the flat projection of the ventral areas, showing the difference in functional correlation between the left and right seeds (voxels more connected with the left seed appear in red, whereas those more connected with the right seed appear in blue). The grid corresponds to the mask presented in (a). e, h, k) The differences between left and right functional correlation values are plotted as a function of the x coordinate. Dots over the curve mark the spots where the difference in connectivity is significantly different ($p < 0.05$, Bonferroni corrected).

Supplementary Figure 1. **Clustering analyses using Pearson correlation** a) The 4 clusters identified in this analysis are presented on a horizontal slice of the infant template. b) Pearson correlation maps showing whole brain connectivity specifically related to each of the 3 defined clusters.

REFERENCES

- Amalric, M., & Dehaene, S. (2016). Origins of the brain networks for advanced mathematics in expert mathematicians. *Proc Natl Acad Sci U S A*, 113(18), 4909-4917.
- Arichi, T., Moraux, A., Melendez, A., Doria, V., Groppo, M., Merchant, N., . . . Edwards, A. D. (2010). Somatosensory cortical activation identified by functional MRI in preterm and term infants. *Neuroimage*, 49(3), 2063-2071.
- Bartocci, M., Bergqvist, L. L., Lagercrantz, H., & Anand, K. J. (2006). Pain activates cortical areas in the preterm newborn brain. *Pain*, 122(1-2), 109-117.
- Biagi, L., Crespi, S. A., Tosetti, M., & Morrone, M. C. (2015). BOLD Response Selective to Flow-Motion in Very Young Infants. *PLoS Biol*, 13(9), e1002260.
- Cai, Q., Paulignan, Y., Brysbaert, M., Ibarrola, D., & Nazir, T. A. (2010). The left ventral occipito-temporal response to words depends on language lateralization but not on visual familiarity. *Cereb Cortex*, 20(5), 1153-1163.
- Caspers, J., Palomero-Gallagher, N., Caspers, S., Schleicher, A., Amunts, K., & Zilles, K. (2015). Receptor architecture of visual areas in the face and word-form recognition region of the posterior fusiform gyrus. *Brain Struct Funct*, 220(1), 205-219.
- Deen, B., Richardson, H., Dilks, D. D., Takahashi, A., Keil, B., Wald, L. L., . . . Saxe, R. (2017). Organization of high-level visual cortex in human infants. *Nat Commun*, 8, 13995.
- Doria, V., Beckmann, C. F., Arichi, T., Merchant, N., Groppo, M., Turkheimer, F. E., . . . Edwards, A. D. (2010). Emergence of resting state networks in the preterm human brain. *Proc Natl Acad Sci U S A*, 107(46), 20015-20020.
- Elman, J. L., Bates, E., Johnson, M. H., Karmiloff-Smith, A., Parisi, D., & Plunkett, K. (1996). *Rethinking innateness: A connectionist perspective on development*. Cambridge, Mass.: MIT Press.
- Fischl, B., Rajendran, N., Busa, E., Augustinack, J., Hinds, O., Yeo, B. T., . . . Zilles, K. (2008). Cortical folding patterns and predicting cytoarchitecture. *Cereb Cortex*, 18(8), 1973-1980.

- Fransson, P., Aden, U., Blennow, M., & Lagercrantz, H. (2011). The functional architecture of the infant brain as revealed by resting-state fMRI. *Cereb Cortex*, 21(1), 145-154.
- Fransson, P., Skiold, B., Engstrom, M., Hallberg, B., Mosskin, M., Aden, U., . . . Blennow, M. (2009). Spontaneous brain activity in the newborn brain during natural sleep--an fMRI study in infants born at full term. *Pediatr Res*, 66(3), 301-305.
- Fransson, P., Skiold, B., Horsch, S., Nordell, A., Blennow, M., Lagercrantz, H., & Aden, U. (2007). Resting-state networks in the infant brain. *Proc Natl Acad Sci U S A*, 104(39), 15531-15536.
- Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., . . . Van Essen, D. C. (2016). A multi-modal parcellation of human cerebral cortex. *Nature*, 536(7615), 171-178.
- Gomez, J., Barnett, M. A., Natu, V., Mezer, A., Palomero-Gallagher, N., Weiner, K. S., . . . Grill-Spector, K. (2017). Microstructural proliferation in human cortex is coupled with the development of face processing. *Science*, 355(6320), 68-71.
- Hannagan, T., Amedi, A., Cohen, L., Dehaene-Lambertz, G., & Dehaene, S. (2015). Origins of the specialization for letters and numbers in ventral occipitotemporal cortex. *Trends Cogn Sci*, 19(7), 374-382.
- Johnson, M. H. (2011). Interactive specialization: a domain-general framework for human functional brain development? *Dev Cogn Neurosci*, 1(1), 7-21.
- Kabdebon, C., Leroy, F., Simmonet, H., Perrot, M., Dubois, J., & Dehaene-Lambertz, G. (2014). Anatomical correlations of the international 10-20 sensor placement system in infants. *Neuroimage*, 99, 342-356.
- Mahmoudzadeh, M., Dehaene-Lambertz, G., Fournier, M., Kongolo, G., Goudjil, S., Dubois, J., . . . Wallois, F. (2013). Syllabic discrimination in premature human infants prior to complete formation of cortical layers. *Proc Natl Acad Sci U S A*, 110(12), 4846-4851.
- Mahmoudzadeh, M., Wallois, F., Kongolo, G., Goudjil, S., & Dehaene-Lambertz, G. (2016). Functional maps at the onset of auditory inputs in very early preterm human neonates. *Cerebral Cortex*.
- Pinel, P., & Dehaene, S. (2009). Beyond hemispheric dominance: brain regions underlying the joint lateralization of language and arithmetic to the left hemisphere. *Journal of Cognitive Neuroscience*, 22(1), 48-66.
- Pinel, P., Lalanne, C., Bourgeron, T., Fauchereau, F., Poupon, C., Artiges, E., . . . Dehaene, S. (2014). Genetic and Environmental Influences on the Visual Word Form and Fusiform Face Areas. *Cereb Cortex*.
- Power, J. D., Schlaggar, B. L., & Petersen, S. E. (2015). Recent progress and outstanding issues in motion correction in resting state fMRI. *Neuroimage*, 105, 536-551.
- Quartz, S. R., & Sejnowski, T. J. (1997). The neural basis of cognitive development: a constructivist manifesto. *Behav Brain Sci*, 20(4), 537-556; discussion 556-596.
- Raichle, M. E. (2015). The restless brain: how intrinsic activity organizes brain function. *Philos Trans R Soc Lond B Biol Sci*, 370(1668).
- Rakic, P. (1988). Specification of cerebral cortical areas. *Science*, 241(4862), 170-176.
- Saygin, Z. M., Osher, D. E., Koldewyn, K., Reynolds, G., Gabrieli, J. D., & Saxe, R. R. (2012). Anatomical connectivity patterns predict face selectivity in the fusiform gyrus. *Nat Neurosci*, 15(2), 321-327.
- Saygin, Z. M., Osher, D. E., Norton, E. S., Youssoufian, D. A., Beach, S. D., Feather, J., . . . Kanwisher, N. (2016). Connectivity precedes function in the development of the visual word form area. *Nat Neurosci*, 19(9), 1250-1255.
- Shum, J., Hermes, D., Foster, B. L., Dastjerdi, M., Rangarajan, V., Winawer, J., . . . Parvizi, J. (2013). A brain area for visual numerals. *J Neurosci*, 33(16), 6709-6715.
- Smith, S. M., Vidaurre, D., Beckmann, C. F., Glasser, M. F., Jenkinson, M., Miller, K. L., . . . Van Essen, D. C. (2013). Functional connectomics from resting-state fMRI. *Trends Cogn Sci*, 17(12), 666-682.

- Smyser, C. D., Inder, T. E., Shimony, J. S., Hill, J. E., Degnan, A. J., Snyder, A. Z., & Neil, J. J. (2010). Longitudinal analysis of neural network development in preterm infants. *Cereb Cortex*, 20(12), 2852-2862.
- Smyser, C. D., Snyder, A. Z., & Neil, J. J. (2011). Functional connectivity MRI in infants: Exploration of the functional organization of the developing brain. *Neuroimage*.
- Toulmin, H., Beckmann, C. F., O'Muircheartaigh, J., Ball, G., Nongena, P., Makropoulos, A., . . . Edwards, A. D. (2015). Specialization and integration of functional thalamocortical connectivity in the human infant. *Proc Natl Acad Sci U S A*, 112(20), 6485-6490.
- van den Heuvel, M. P., Kersbergen, K. J., de Reus, M. A., Keunen, K., Kahn, R. S., Groenendaal, F., . . . Benders, M. J. (2015). The Neonatal Connectome During Preterm Brain Development. *Cereb Cortex*, 25(9), 3000-3013.
- Weiner, K. S., Barnett, M. A., Lorenz, S., Caspers, J., Stigliani, A., Amunts, K., . . . Grill-Spector, K. (2016). The Cytoarchitecture of Domain-specific Regions in Human High-level Visual Cortex. *Cereb Cortex*.
- Weiner, K. S., Golarai, G., Caspers, J., Chuapoco, M. R., Mohlberg, H., Zilles, K., . . . Grill-Spector, K. (2014). The mid-fusiform sulcus: a landmark identifying both cytoarchitectonic and functional divisions of human ventral temporal cortex. *Neuroimage*, 84, 453-465.